

Lysine Acetylsalicylate for Injection

For IV use

COMPOSITION :

Lysine Acetylsalicylate for Injection 0.5g

Each vial contains:

Lysine Acetylsalicylate 0.9 g
eq. to Acetylsalicylic Acid BP..... 0.5 g
Glycine BP..... 0.1 g

Lysine Acetylsalicylate for Injection 1g

Each vial contains:

Lysine Acetylsalicylate 1.8 g
eq. to Acetylsalicylic Acid BP 1g
Glycine BP..... 0.2 g

PHARMACEUTICAL FORM : Powder for Injection

INDICATIONS:

Moderate and severe pain of various origins, rheumatic pain, hyperthermia, thromboembolic events, heart attack, stroke.

DOSAGE AND METHOD OF ADMINISTRATION:

Lysine Acetylsalicylate for Injection 0.5g

Adults: 1 or 2 vials per injection; for severe pain: 2 vials in a single I.V. injection. Usual 24 hour dosage is 1 to 4 vials. In some cases, daily dosage can be increased to 8 vials.

Children: 10-25 mg/kg/day. For children from 6-10 years: 1/2 vial per day. Older than 11 years: 1-2 vials per day.

Lysine Acetylsalicylate for Injection 1 g

Adults: Usual dosage 1 vial per injection twice daily. Maximal dosage: 4 vials per day.

Or as directed by the Physician.

MODE OF ADMINISTRATION: For IV use

DIRECTION FOR USE: For 0.5g Dissolve the contents of vial in 5ml of water for injection.

For 1g Dissolve the contents of vial in 10ml of water for injection.

CONTRAINDICATIONS:

Gastroduodenal ulcer, history of hypersensitivity to salicylates; any hemorrhagic disorders (hereditary or acquired); any risk of hemorrhage; last quarter of pregnancy. Use is not recommended in: gout, menorrhagia.

DRUG INTERACTIONS:

Vitamin K blockers and Heparin, Sulfonylurea, Methotrexate, NSAIDs, Uricosuric agents.

PRECAUTIONS:

Use with caution in case of: associated anti-coagulant therapy, asthma, renal insufficiency; first two quarters of pregnancy, IUD.

SIDE EFFECTS :

Side-effects are those of acetylsalicylic acid. The more common side effects of acetylsalicylic acid include:

- Hypersensitivity (which may mean you have skin rashes or itching, or wheezing or coughing or difficulty breathing)
- Nausea & vomiting
- Ringing in your ears
- Pain or discomfort in your stomach
- Acetylsalicylic acid may sometimes cause asthma or bronchospasm (narrowing of the airways that causes wheezing or difficulty breathing).
- Acetylsalicylic acid may cause bleeding, and you must tell your doctor if you experience any unusual bleeding.

STORAGE:

Store at temperature below 25°C. Protect from light & moisture. Do not freeze.

KEEP MEDICINE OUT OF REACH OF CHILDREN.

PRESENTATION :

50 vials are packed in a printed carton.

Commercialisé Par:
Kwality Pharmaceuticals Ltd.
INDIA.