


PRAZIQUANTEL TABLETS U.S.P.

DISTOSIDE® Tablets 600mg

DESCRIPTION : Distoside (Praziquantel) is 4H-Pyrazino[2,1-a]isoquinolin-4-one, 2-cyclohexylcarbonyl-1,2,3,6,7,11b-hexahydro2-(Cyclohexylcarbonyl)-1,2,3,6,7,11b-hexahydro-4H-pyrazino-[2,1-a]isoquinoline-4-one—[55268-74-1].



M. W. : 312.41

Empirical formula: C₁₈ H₂₄ N₂ O₂

It is colourless and almost odourless crystalline compound with bitter taste.

COMPOSITION : Each film coated tablet contains

Praziquantel U.S.P. 600mg

Excipients q.s.

Colour - Titanium Dioxide

INDICATIONS : Distoside is effective to all Schistosome species pathogenic to man including wide range of Cestodes and liver and lung flukes.

DOSAGE : Dosage must be adjusted to the individual case by the attending physician. According to the result obtained so far, the following dosages are recommended.


Dosage Regimen

Parasites	Dosage
Schistosoma haematobium	20mg/kg x 2 for 1 day
S. Mansoni	40mg/kg x 1 for 1 day
S. Japonicum	20mg/kg x 3 for 1 day
Clonorchis sinensis	25mg/kg x 3 for 1 day
Opisthorchis viverrini	25mg/kg x 3 for 1 day
Paragonimus westermani	25mg/kg x 3 for 2 days
P. Heterotremus	25mg/kg x 3 for 2 days

The numbers of tablets required for one single dose can be calculated from the following tablets.

Body weight in (kg)	No. Of tablets (25mg/kg)	Body weight in (kg)	No. Of tablets (20mg/kg)
22 - 26	1	20 - 25/26	3/4
27 - 33	1 1/4	26/27 - 33	1
34 - 38	1 1/2	34 - 41	1 1/4
39 - 44	1 3/4	42 - 48	1 1/2
45 - 50	2	49 - 56	1 3/4
51 - 56	2 1/4	57 - 63	2
57 - 62	2 1/2	64 - 70	2 1/4
63 - 68	2 3/4	71 - 78	2 1/2
69 - 75	3	79 - 86	2 3/4

ADMINISTRATION : The tablets should be washed down at once unchewed with some liquid best after a meal. In case of repeated intake on the same day, the interval between the individual doses should not be less than 4 and not more than 6 hours. One oblong tablets has three scores. Each segment contains 150mg active ingredient so that dosage can be exactly adjusted to the patient's body weight. If 1/4 of a tablet is required, this is best achieved by breaking the segment from the outer end. Segments are broken off by pressing notch with thumb nails.



CONTRAINDICATIONS : None known so far.

PRECAUTIONS : 1. Side Effects : It is mild, transient in nature and tolerable.

Dizziness, headache, drowsiness, nausea, abdominal discomfort, urticaria, rise in temperature may or may not occur.

2. General Precaution : Side effects are likely to occur more often after single dose of 40mg/kg and/or in case of a patients with heavy worm burden. In patients with liver impairment, no additional adverse effects have occurred. For the treatment of cerebral and ocular cysticercosis, it is necessary to hospitalize the patient and to secure the attendance of an expert neurologist.

3. During pregnancy and breast-feeding;

* In accordance with the general rules governing the use of drugs during pregnancy, Distoside should not be taken during the first three months of pregnancy, although animal experiments do not point out to harmful effects on either pregnant mother animal or unborn baby.

* A breast-feeding mother can undergo treatment provided her baby is fed otherwise on the day of treatment and during the subsequent 72 hours.

PACKAGE : 2 x 4 tablets

STORAGE : Store below 25°C.

Protect from light & moisture.

Manufactured by :

Chandra Bhagat Pharma Pvt. Ltd.

103, Shivam Complex, Library Road, Navsari.

Regd. Off.: 323-F, Bhagat Bhuvan,

Dr. Ambedkar Road, Matunga (E),

Mumbai - 400 018.

At: Old National Highway No. 8,

Sankarda - 391 350,

Vadodra, Gujarat.

Email: info@cbpharma.com

Site : www.cbpharma.com