

Pyzina 500 MG Tablet

Manufacturer: Lupin Ltd

Medicine composition: [Pyrazinamide](#)

Prescription vs.OTC: Prescription by Doctor required

Last Updated: February 04, 2020

About Pyzina 500 MG Tablet

Pyzina 500 MG Tablet is an anti-tubercular medicine, which is used alongside other medicines to treat tuberculosis. It also prevents the infection from spreading. If you notice some unwanted effects like a [fever](#), tenderness in the abdomen, [jaundice](#), nausea, vomiting or lack of appetite inform your doctor immediately.

This medicine is not recommended during pregnancy or breastfeeding unless absolutely necessary. Consult your doctor about the potential risks and benefits before starting your treatment with this medicine.

If you suffer from the medical conditions mentioned here, inform your doctor so extra care can be taken during the treatment:

[pregnant](#) or breast-feeding.

kidney or [liver problems](#), alcoholism, acute gout, [diabetes](#) or [porphyria](#).

If you are taking any other medicines including those which are available without a prescription as well as herbal and complementary medicines.

Allergic to any medicine.

This medicine will be prescribed to you and you will be asked to take it three times a week during the first two months of your treatment under the supervision of a healthcare provider. Taking the tablets regularly and on time is very important.

It is also important to complete the course of treatment so that there are no chances of the infection to come back. The effect of this medicine can be observed within the first 2 hours of taking it and lasts for an average duration of 27 to 30 hours.

Information given here is based on the salt content of the medicine. Uses and effects of the medicine may vary from person to person. It is advisable to consult a [Pulmonologist](#) before using this medicine.

What are the uses of Pyzina 500 MG Tablet

The medication can treat the below provided problems:

Tuberculosis

This medication is used in the treatment of tuberculosis which is a lung disease caused by [Mycobacterium](#) tuberculosis.

What are the contraindications of Pyzina 500 MG Tablet ?

[Allergy](#)

Not recommended in patients with known allergy to this medication.

Severe Liver Impairment

Not recommended in the patients with severe liver impairment.

Acute Gout

Not recommended in patients with increased uric acid levels or acute gout.

[Porphyria](#)

What are the side effects of Pyzina 500 MG Tablet ?

A patient can suffer from the following side effects:

[Joint Pain](#)

Loss Of Appetite

Unusual Tiredness And Weakness

Yellow Colored Eyes Or Skin

[Itching](#)

[Skin Rash](#)

[Anorexia](#)

Flushing

Pruritus

Urticaria

Photosensitivity

Key highlights of Pyzina 500 MG Tablet

How long is the duration of effect?

The effect of this medicine lasts for an average duration of 27 to 30 hours.

What is the onset of action?

The effect of this medicine can be observed in 2 hours.

Are there any pregnancy warnings?

This medicine is not recommended for use during pregnancy unless absolutely necessary.

Is it habit forming?

No habit forming tendencies were reported.

Are there any breast-feeding warnings?

This medicine is excreted in breast milk. It is recommended for breastfeeding women only if clearly needed.

What are the dosage instructions?

Missed Dose instructions

If you miss a dose of medication, take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose. Do not double your dose to make up for the missed dose.

Overdose instructions

Seek emergency medical treatment or contact the doctor in case of an overdose.

Where is the Pyzina 500 MG Tablet approved?

India

United States

Japan

How does this medication work?

This medicine belongs to an antitubercular class of drugs. It works by inhibiting the growth of the organism by decreasing the pH inside the cell by converting it into an active metabolite called pyrazinoic acid in the presence of pyrazinamidase.

What are the interactions of Pyzina 500 MG Tablet ?

Whenever you take more than one medicine, or mix it with certain foods or beverages, you're at risk of a drug interaction.

Interaction with Alcohol

Alcohol

Interaction with alcohol is unknown. It is advisable to consult your doctor before consumption.

Interaction with Medicine

Methotrexate

These medicines may increase the risk of liver injury. Inform the doctor if you are receiving these medicines. The baseline of liver function tests should be obtained before starting the treatment. Any signs and symptoms of elevated liver enzymes, yellow colored eyes, and skin, dark urine should be reported to the doctor.

Cholera Vaccine

If you have taken this medication, then it is advisable to wait for 14 days before taking the Cholera vaccine. Usage of other antibiotics and vaccines should be reported to the doctor.

Interaction with Disease

Gout

This medicine is not recommended in patients with gout. Inform the doctor about your medical condition. Serum uric acid levels should be monitored carefully. Discontinue the treatment if the symptoms persist.

Interaction with Food

Food

Information not available.

FAQs about Pyzina 500 MG Tablet

Ques: What is Pyzina 500 MG Tablet?

Ans: is an anti-tuberculosis medicine which is used alongside other medicines to treat tuberculosis.

Ques: How long do I need to use Pyzina 500 MG Tablet before I see improvement in my condition?

Ans: This medication is to be taken till the time you see an improvement in your health conditions.

Ques: At what frequency do I need to use Pyzina 500 MG Tablet?

Ans: This medication will be prescribed to you and you will be asked to take it three times a week, during the first two months of your treatment

Ques: Should I use Pyzina 500 MG Tablet empty stomach, before food or after food?

Ans: This medication should be taken after food in a prescribed dosage.

Ques: What are the instructions for the storage and disposal of Pyzina 500 MG Tablet?

Ans: This medication should be kept in a cool dry place and in its original packaging. Make sure this medication remains unreachable to children and pets.



References

Pyrazinamide- Drug Information Portal [Internet]. druginfo.nlm.nih.gov. 2019 [Cited 11 December 2019]. Available from:

<https://druginfo.nlm.nih.gov/drugportal/name/pyrazinamide>

Pyrazinamide- DrugBank [Internet]. Drugbank.ca. 2019 [Cited 11 December 2019]. Available from:

<https://www.drugbank.ca/drugs/DB00339>

Zinamide 500mg Tablets- EMC [Internet] medicines.org.uk. 2015 [Cited 11 December 2019].

Available from:

<https://www.medicines.org.uk/emc/product/5273/smpc>